arXiv:1110.5038v3 [math.CO] 6 Jul 2012

Lifting automorphisms of abelian regular coverings of graphs

Haimiao Chen *

Department of mathematics, Peking University, Beijing, China

Abstract

This article proposes an effective criterion for lifting automorphisms of regular coverings of graphs, with the covering transformation group being any finite abelian group.

1 Introduction

First let us recall some terminologies in graph theory. For more one can refer to [2], [3], [4], [7], [9].

We suppose that all graphs are finite, simple and undirected. For a graph Γ, denote an edge connecting the vertices u, v by $\{u, v\}$; each edge $\{u, v\}$ gives rise to a pair of opposite arcs (u, v) and (v, u). Use $V(\Gamma), E(\Gamma), A(\Gamma)$ to denote the set of vertices, edges, arcs of Γ, respectively, and use $Aut(\Gamma)$ for the automorphism group.

A graph is the same thing as a one-dimensional simplicial complex (see [5], page 104) with orientation forgotten. A covering of graphs $\pi : \tilde{\Gamma} \to \Gamma$ is a simplicial covering map of one-dimensional simplicial complexes. It is called regular (or an A-covering) if there is a subgroup A of the covering transformation group K that acts regularly on each fiber. Moreover $A = K$ if Γ is connected.

An automorphism $\alpha \in Aut(\Gamma)$ is lifted to $\tilde{\alpha} \in Aut(\tilde{\Gamma})$ if $\pi\tilde{\alpha} = \alpha\pi$. The image of an arc h (or a walk W) is denoted h^{α} (respectively W^{α}).

---

*Email: *chenhm@math.pku.edu.cn*

Let A be a finite group. An A-voltage assignment of Γ is a function $\phi : A(\Gamma) \to A$ such that $\phi(u,v) = \phi(v,u)^{-1}$, for all $(u,v) \in A(\Gamma)$; the values of ϕ are called voltages. The graph $\tilde{\Gamma} = \Gamma \times_\phi A$ derived from ϕ is defined by $V(\tilde{\Gamma}) = V(\Gamma) \times A$, and $E(\tilde{\Gamma}) = \{\{(u,g),(v,\phi(u,v)g)\} \colon \{u,v\} \in E(\Gamma), g \in A\}$. The first coordinate projection $\pi : \tilde{\Gamma} = \Gamma \times_\phi A \to \Gamma$ makes an A-coverings.

Given a spanning tree T of Γ, each edge $e \in E(\Gamma) - E(T)$ is called cotree. A voltage assignment ϕ is called T-reduced if the voltage associated to each tree arc is the identity. It was shown in [3] that every regular covering of Γ is isomorphic to a derived graph from a T-reduced voltage assignment with respect to an arbitrary tree T.

The voltage assignment ϕ naturally extends to walks in Γ. For any walk W, let $\phi(W)$ denote the voltage of W.

We are able to give a nice solution to the following problem:

Problem 1.1. *Given a graph Γ, a regular covering $\Gamma \times_\phi A$ with A a finite abelian group, and an automorphisms α of Γ, decide whether α can be lifted.*

The following proposition of [7] plays a key role:

Proposition 1.2. *Let $\tilde{\Gamma} = \Gamma \times_\phi A$ be a regular covering. Then an automorphism α of Γ can be lifted to an automorphism of $\tilde{\Gamma}$ if and only if, for each closed walk W in Γ, one has $\phi(W^\alpha) = 0$ if $\phi(W) = 0$.*

Recently much attention has been paid to the automorphism lifting problem. The motivation was to construct highly transitive graphs, and now is from different sources, for instance, to classify symmetries of graphs. In [2] a linear criteria is proposed for the case that A is elementary abelian, and applied to classify arc-transitive A-coverings of the Petersen graph. In [9] the authors classify all vertex-transitive elementary abelian coverings of the Petersen graph; the method, developed by themselves in [8], is finding invariant subspaces of a group representation. Other results are [6], [10], and so on.

To the automorphism lifting problem for arbitrary finite abelian groups, we contribute a criterion based on a theory for diagonalizing matrices over the ring $\mathbb{Z}/p^k$. The criterion is easy to run, in that, one only needs to calculate some matrices derived from input data, and check whether the entries of the resulted matrices satisfy certain conditions. We believe that classifying finite abelian symmetries of graphs, at least for small graphs, is possible in the near future.

We discuss a property of matrices over $\mathbb{Z}/p^k$ in Section 2, and apply it to give a criterion for lifting automorphisms in Section 3; in Section 4 a practical algorithm is presented and illustrated via an example.

Here are some notational conventions. Operations in abelian groups are written additively. For a matrix C, denote its (i,j)-entry by $C_{i,j}$, and we often write $C = (C_{i,j})$. For a ring $\mathcal{R}$, let $\mathcal{M}_{m\times n}(\mathcal{R})$ be the set of all $m \times n$-matrices with entries in $\mathcal{R}$. Let $\mathcal{M}_n(\mathcal{R}) = \mathcal{M}_{n\times n}(\mathcal{R})$, and let $GL(n, \mathcal{R})$ be the set of invertible ones.

2 On matrices over $\mathbb{Z}/p^k$

Let p be a prime number, and $R = \mathbb{Z}/p^k$. The ring R is no longer an integral domain. Nevertheless, we shall see that finitely generated modules over R share some structures with those over a Euclidean domain.

We can talk about the divisibility of an element $\lambda \in R$ by $p^r, 0 \leqslant r < k$, without ambiguity. Note that λ is invertible if and only if it is not divisible by p; it follows that if λ is invertible and λ' is not, then $\lambda + \lambda'$ is invertible.

Also, for $X \in \mathcal{M}_{m\times n}(R), K \in \mathcal{M}_{k\times l}(R), C \in \mathcal{M}_{n\times k}(\mathbb{Z})$, it is meaningful to talk about $XC \in \mathcal{M}_{m\times k}(R), CK \in \mathcal{M}_{n\times l}(R)$.

Definition 2.1. *For $0 \neq \lambda \in R$, the* p-degree *of λ, denoted $d_p(\lambda)$, is the largest integer $r, 0 \leqslant r < k$ such that $p^r|\lambda$. For convention set $d_p(0) = k$.*

Definition 2.2. *A nonzero matrix $X = (X_{i,j}) \in \mathcal{M}_{m\times n}$ is* in normal form *if $X_{i,j} = \delta_{i,j}p^{r_i}$ for some integers $r_1, \cdots, r_m$, with $0 \leqslant r_1 \leqslant \cdots \leqslant r_l < k = r_{l+1} = \cdots = r_m$ for some $l \leqslant min\{m, n\}$.*

Lemma 2.3. *For each $0 \neq X \in \mathcal{M}_{m\times n}(R)$, there exist $Q \in GL(m, R), T \in GL(n, R)$ such that QXT is in normal form.*

Proof. The result is analogous to that of matrices with entries in $\mathbb{Z}$ (see [1], page), and is proved by showing that X can be changed into normal form by a series of row and column transformations.

Choose an entry $X_{i_1,j_1} \neq 0$ with smallest p-degree, and let $r_1 = d_p(X_{i_1,j_1})$. Interchange the i_1-th row of X with the first row, and the j_1-th column with the first column, and divide the first row by $X_{i_1,j_1}/p^{r_1}$. The matrix obtained has $(1,1)$-entry p^{r_1}, dividing all entries. Then perform row-transformations to eliminate the $(i,1)$-entries for $1 < i \leqslant m$. Denote the new matrix by $X^{(1)} = (X^{(1)}_{i,j})$.

Doing the same thing to the submatrix $(X^{(1)}_{i+1,j+1}), 1 \leqslant i < m, j < n$, of $X^{(1)}$ changes $X^{(1)}$ into $X^{(2)} = (X^{(2)}_{i,j})$, with $X^{(2)}_{1,1} = p^{r_1}$, $X^{(2)}_{2,2} = p^{r_2}$ dividing $X^{(2)}_{i,j}, \forall i \geqslant 2$; $r_1 \leqslant r_2$, and $X^{(2)}_{i,j} = 0$ for $i > j, j \in \{1, 2\}$.

Continue this process. Eventually there is some $l \leqslant m, n$, such that $X^{(l)} = (X^{(l)}_{i,j})$ satisfies $X^{(l)}_{i,i} = p^{r_i} | X^{(l)}_{i,j}, \forall 1 \leqslant i \leqslant l$; $r_1 \leqslant \cdots \leqslant r_l < k$; $X^{(l)}_{i,j} = 0$ for $j < i \leqslant l$ or $i > l$.

Finally, perform column transformations to $X^{(l)}$, to eliminate all the "off-diagonal" entries. Then the resulted matrix is in normal form. □

The following corollary, which is an analogue of the structure theorem of finitely modules over a Euclidean domain (see [1] page 475), is easy to prove. We state it without proof since it is never applied in this article.

Corollary 2.4. *Every finitely generated $\mathbb{Z}/p^k$-module has a set of generators $\{x_1, \cdots, x_l\}$ such that there is no relation other than $p^{r_i} x_i = 0, i \in \{1, \cdots, l\}$, with $0 < r_1 \leqslant \cdots \leqslant r_l < k$ for some integers $r_1, \cdots, r_l$.*

In the end of this section, we point out that all the results here can be generalized when $\mathbb{Z}/p^k$ is replaced by $\mathbb{Z}/N$ for any natural number N.

3 Criteria for lifting automorphisms

3.1 Set up

From now on, we make the followings assumptions.

(1) The abelian group $A = \prod_{\gamma=1}^{g} A_\gamma$, with $A_\gamma = \prod_{\eta=1}^{a_\gamma} \mathbb{Z}/p_\gamma^{k(\gamma,\eta)}$, the prime numbers $p_1, \cdots, p_g$ being distinct, and $k(\gamma, 1) \leqslant \cdots \leqslant k(\gamma, a_\gamma)$ for each γ.

(2) For each edge $e \in E(\Gamma)$, one of its arcs is chosen and denoted $h(e)$.

(3) A spanning tree T and a vertex $v_0 \in V(T)$ are chosen. For $v_1, v_2 \in V(\Gamma) = V(T)$, let $W(v_1, v_2)$ denote the unique reduced walk from v_1 to v_2. Let $e_1, \cdots, e_t$ be all the cotree edges, and for $1 \leqslant i \leqslant t$, suppose the chosen arc for e_i is $h(e_i) = (u_i, v_i)$; let L_i be the unique closed reduced walk based at v_0 and containing $h(e_i)$, that is, $L_i = W(v_0, u_i) \cup h(e_i) \cup W(v_i, v_0)$.

(4) The voltage assignment ϕ is T-reduced, so $\phi(h(e)) = 0, \forall e \in E(T)$.

It is well-known that the first homology group $H_1(\Gamma; \mathbb{Z}) \cong \mathbb{Z}^t$ (see [5], page 43) has a basis represented by $L_1, \cdots, L_t$.

The voltage assignment ϕ induces a group homomorphism

$$\overline{\phi}: H_1(\Gamma; \mathbb{Z}) \to A. \tag{1}$$

Now each automorphism $\alpha \in Aut(\Gamma)$ induces an isomorphism

$$\alpha_*: H_1(\Gamma; \mathbb{Z}) \to H_1(\Gamma; \mathbb{Z}). \tag{2}$$

Let $S^\alpha \in GL(t, \mathbb{Z})$ denote the corresponding matrix, so that

$$\alpha_*(L_i) = \sum_{j=1}^{t} (S^\alpha)_{i,j} L_j. \tag{3}$$

Proposition 1.2 can be reformulated as

Proposition 3.1. *The automorphism α can be lifted if and only if*

$$Ker(\overline{\phi}) = Ker(\overline{\phi} \circ \alpha_*). \tag{4}$$

Denote $\theta_i = \overline{\phi}(L_i) = \phi(L_i) \in A$. Suppose $\theta_i = (\theta_i^{(1)}, \cdots, \theta_i^{(g)})$, with $\theta_i^{(\gamma)} \in A_\gamma$, and for each γ, suppose further that $\theta_i^{(\gamma)} = (\theta_i^{(\gamma)}(1), \cdots, \theta_i^{(\gamma)}(a_\gamma))$, with $\theta_i^{(\gamma)}(\eta) \in \mathbb{Z}/p_\gamma^{k(\gamma,\eta)}$.

Note that for all $1 \leqslant \eta \leqslant a_\gamma$ there is a monomorphism

$$\iota_{\gamma,\eta}: \mathbb{Z}/p_\gamma^{k(\gamma,\eta)} \hookrightarrow \mathbb{Z}/p_\gamma^{k(\gamma,a_\gamma)}, \quad \lambda \mapsto p_\gamma^{k(\gamma,a_\gamma)-k(\gamma,\eta)} \cdot \lambda \tag{5}$$

Let $B_\gamma \in \mathcal{M}_{t\times a_\gamma}(\mathbb{Z}/p_\gamma^{k(\gamma,a_\gamma)})$ denote the matrix given by

$$(B_\gamma)_{i,j} = \iota_{\gamma,j}(\theta_i^{(\gamma)}(j)). \tag{6}$$

Then $w = (w_1, \cdots, w_t) \in \mathbb{Z}^t \in Ker(\overline{\phi})$ is equivalent to

$$wB_\gamma = 0, \quad \forall 1 \leqslant \gamma \leqslant g. \tag{7}$$

At last, to be convenient, we make the following

Definition 3.2. *For a ring $\mathcal{R}$, and a matrix $X \in \mathcal{M}_{m\times n}(\mathcal{R})$, the* zero set *of X* in *$\mathcal{R}^m$, denoted $Z(X)$, is the set of the row vectors $w = (w_1, \cdots, w_m) \in \mathcal{R}^m$ satisfying $wX = 0$, or explicitly, $\sum_{i=1}^{m} w_i X_{i,j} = 0, \forall 1 \leqslant j \leqslant n$.*

3.2 The case of one prime factor

Suppose $g = 1$. We simplify notations by $p = p_1$, $a = a_1$, $k = k(1, a)$, $B = B_1$, $R = \mathbb{Z}/p^k$.

For a matrix $X \in \mathcal{M}_{l\times m}(R)$, let $\langle X\rangle$ denote the subgroup of R^m generated by the row vectors of X.

Since $B \in \mathcal{M}_{t\times a}(R)$, by Lemma 2.3 one can choose $Q \in GL(t, R), T \in GL(a, R)$ so that $QBT = B^0$ is in normal form, $(B^0)_{i,j} = \delta_{i,j}p^{s_i}$.

Let $P \in \mathcal{M}_{(t-i_0+1)\times t}(R)$ with $P_{i,j} = \delta_{i+i_0-1,j} \cdot p^{k-s_j}$. Clearly $Z(B^0) = \langle P\rangle$ and $Z(B) = Z(B^0)Q = \langle PQ\rangle$.

Theorem 3.3. *Choose Q, T such that $B^0 = QBT$ is in normal form, $(B^0)_{i,j} = \delta_{i,j}p^{s_i}$, with i_0 the smallest i such that $s_i > 0$. Then α can be lifted if and only if* $\deg_p((QS^\alpha Q^{-1})_{i,j}) \geqslant s_i - s_j, \forall i > j$.

Proof. By Proposition 3.1, α can be lifted if and only if $Z(S^\alpha B) = Z(B)$, that is, $\langle PQS^\alpha\rangle = \langle PQ\rangle$.

If $\langle PQS^\alpha\rangle = \langle PQ\rangle$, then $P(QS^\alpha Q^{-1}) = UP$ for some $U \in \mathcal{M}_{t-i_0+1}(R)$, implying $\deg_p((QS^\alpha Q^{-1})_{i,j}) \geqslant s_i - s_j, \forall i > j$.

Conversely, if $\deg_p((QS^\alpha Q^{-1})_{i,j}) \geqslant s_i - s_j, \forall i > j$, then, using the characterization that $(w_1, \cdots, w_t) \in \langle P\rangle$, it is easy to see that $\langle P(QS^\alpha Q^{-1})\rangle \leqslant \langle P\rangle$. Since the subgroup $\langle P(QS^\alpha Q^{-1})\rangle$ has as many elements as $\langle P\rangle$ does, we have $\langle P(QS^\alpha Q^{-1})\rangle = \langle P\rangle$, and hence $\langle PQS^\alpha\rangle = \langle PQ\rangle$.

□

3.3 The general case

Let $N = \prod_{\gamma=1}^{g} p_\gamma^{k(\gamma,a_\gamma)}$, $R_\gamma = \mathbb{Z}/p_\gamma^{k(\gamma,a_\gamma)}$ for $\gamma \in \{1, \cdots, g\}$, and let $R = \mathbb{Z}/N \cong R_1 \times \cdots \times R_g$. There are canonical projections $q_\gamma : R \twoheadrightarrow R_\gamma$ and injections $\iota_\gamma : R_\gamma \hookrightarrow R$ which induce maps $q_\gamma : R^t \twoheadrightarrow R^t_\gamma$ and $\iota_\gamma : R^t_\gamma \hookrightarrow R^t$, respectively.

Let $B \in \mathcal{M}_{t\times(\sum_{\gamma=1}^{g} a_\gamma)}(R)$ be given by, for $\sum_{\eta=1}^{\gamma-1} a_\eta < j \leqslant \sum_{\eta=1}^{\gamma} a_\eta$,

$$B_{i,j} = \iota_\gamma((B_\gamma)_{i,j-\sum_{\eta=1}^{\gamma-1} a_\eta}). \tag{8}$$

Lemma 3.4. *$Z(SB) = Z(B)$ in R^t if and only if $Z(SB_\gamma) = Z(B_\gamma)$ in R^t_γ for each γ.*

Proof. Note that $w \in Z(B)$ if and only if $q_\gamma(w) \in Z(B_\gamma), \forall \gamma$; $w \in Z(SB)$ if and only if $q_\gamma(w) \in Z(SB_\gamma), \forall \gamma$. Moreover, for each γ, a vector $w^{(\gamma)} \in Z(B_\gamma)$ if and only if $\iota_\gamma(w^{(\gamma)}) \in Z(B)$, and $w^{(\gamma)} \in Z(SB_\gamma)$ if and only if $\iota_\gamma(w^{(\gamma)}) \in Z(SB)$.

The lemma then follows. $\square$

Since α can be lifted if and only if $Z(SB) = Z(B)$ in R^t, applying this lemma we are able to establish the following

Theorem 3.5. *For $\gamma \in \{1, \cdots, g\}$, choose $Q_\gamma \in GL(t, R_\gamma), T_\gamma \in GL(a_\gamma, R_\gamma)$ so that $B^0_\gamma = Q_\gamma B_\gamma T_\gamma$ is in normal form, $(B^0_\gamma)_{i,j} = \delta_{i,j} p_\gamma^{s_{\gamma,i}}$, with $i_{\gamma,0}$ the smallest i such that $s_{\gamma,i} > 0$. Then α can be lifted if and only if for each γ, $\deg_{p_\gamma}((Q_\gamma S^\alpha Q_\gamma^{-1})_{i,j}) \geqslant s_{\gamma,i} - s_{\gamma,j}, \forall i > j$.*

4 Conclusion and example

Summarizing the last section, we write down the algorithm.

Algorithm 4.1. *Under the assumptions in the beginning of Section 3.1.*

1. Determine the "basic loops" $L_1, \cdots, L_t$. Let $\theta_i = \phi(L_i) \in A$, make the matrices B_γ for $1 \leqslant \gamma \leqslant g$, as in (6).

2. Calculate the matrix S^α representing the action of α on $H_1(\Gamma; \mathbb{Z})$, that is, $\alpha_*(L_i) = \sum_{j=1}^{t} (S^\alpha)_{i,j} L_j$.

3. For each γ, choose Q_γ, T_γ to normalize B_γ so that $(Q_\gamma B_\gamma T_\gamma)_{i,j} = \delta_{i,j} p_\gamma^{s_{\gamma,i}}$, with $0 = s_{\gamma,1} = \cdots = s_{\gamma,i_{\gamma,0}-1} < s_{\gamma,i_{\gamma,0}} \leqslant \cdots \leqslant s_{\gamma,t}$.

4. If $\deg_{p_\gamma}((Q_\gamma S^\alpha Q_\gamma^{-1})_{i,j}) \geqslant s_{\gamma,i} - s_{\gamma,j}$, $\forall i > j$, $\forall \gamma$, then α can be lifted. Otherwise not.

The matrix S^α can be determined as follows: the entry $S^\alpha_{i,j}$ is the algebraic sum of the times that $h(e_j)$ appears in the walk $(L_i)^\alpha$. Here an arc is denoted $-h(e_j)$ if it is opposite to $h(e_j)$.

We now illustrate this algorithm by an example.

Example 4.2. *Take Γ to be the Petersen graph. Let $X = \{a, b, c, d, e\}$, and $V = \{0, 1, 2, 3, 4, 5, 6, 7, 8, 9\}$, where $0 = \{a, b\}$, $1 = \{c, d\}$, $2 = \{c, e\}$, $3 = \{d, e\}$, $4 = \{a, e\}$, $5 = \{b, e\}$, $6 = \{a, d\}$, $7 = \{b, d\}$, $8 = \{a, c\}$, $9 = \{b, c\}$. The Petersen graph has $V(\Gamma) = V$ and $E(\Gamma) = \{\{i, j\} \colon i, j \in V, i \cap j = \emptyset\}$. It is known that $Aut(\Gamma) \cong S_5$, the action being induced from that on X.*

Fix a spanning tree T *as in Figure 1(a),* $v_0 = 0$*; the induced subgraph is shown in (b). For each cotree edge* $\{i, j\}$*, choose the arc* (i, j) *with* $i < j$*. Let* $h_1 = (5, 8), h_2 = (7, 8), h_3 = (4, 7), h_4 = (4, 9), h_5 = (6, 9), h_6 = (5, 6)$*.*

Take $A = \mathbb{Z}/2 \times \mathbb{Z}/2 \times \mathbb{Z}/4$*.*

Define the T*-reduced voltage assignment* ϕ *by* $\theta_1 = (1, 1, 1)$*,* $\theta_2 = (1, 0, 2)$*,* $\theta_3 = (1, 1, 2)$*,* $\theta_4 = (1, 0, 3)$*,* $\theta_5 = (1, 1, 0)$*,* $\theta_6 = (1, 0, 0)$*.*

Let $\alpha_1 = (0)(2)(13)(67)(49)(58), \quad \alpha_2 = (4)(7)(19)(56)(28)(03),$
$\alpha_3 = (0)(123)(468)(579), \quad \alpha_4 = (0)(1)(2)(3)(45)(67)(89).$



Figure 1: (a) The spanning tree T; (b) the induced subgraph

Now the algorithm is run as followings.

1. We have $t = 6$. From $\theta_1, \cdots, \theta_6$ we obtain $B = \begin{pmatrix} 2 & 2 & 1 \\ 2 & 0 & 2 \\ 2 & 2 & 2 \\ 2 & 0 & 3 \\ 2 & 2 & 0 \\ 2 & 0 & 0 \end{pmatrix}$.

2. Determine the matrices $S_k = S^{\alpha_k}$ for $1 \leqslant k \leqslant 4$:

$$S_1 = \begin{pmatrix} -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \end{pmatrix}, S_2 = \begin{pmatrix} 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & -1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & -1 & -1 \end{pmatrix},$$

$$S_3 = \begin{pmatrix} 0 & 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 \\ -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \end{pmatrix}, S_4 = \begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}.$$

3. Normalize B into $B^0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$ by $T = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}$ and

$$Q = \begin{pmatrix} 3 & 2 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 \\ 2 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 1 & 0 & 1 & 0 \\ 2 & 1 & 0 & 0 & 0 & 1 \end{pmatrix}. \text{ So } Q^{-1} = \begin{pmatrix} 1 & 3 & 3 & 0 & 0 & 0 \\ 2 & 2 & 3 & 0 & 0 & 0 \\ 2 & 3 & 1 & 0 & 0 & 0 \\ 3 & 0 & 1 & 1 & 0 & 0 \\ 0 & 3 & 1 & 0 & 1 & 0 \\ 0 & 0 & 3 & 0 & 0 & 1 \end{pmatrix},$$

and $s_1 = 0, s_2 = s_3 = 1, s_4 = s_5 = s_6 = 2$; $i_0 = 2$.

4. Calculating $QS_kQ^{-1}, k \in \{1, 2, 3, 4\}$, the results are

$$QS_1Q^{-1} = \begin{pmatrix} 1 & 0 & 0 & 0 & 3 & 2 \\ 0 & 1 & 0 & 0 & 3 & 3 \\ 2 & 2 & 3 & 0 & 0 & 3 \\ 0 & 2 & 0 & 3 & 3 & 3 \\ 0 & 0 & 0 & 0 & 3 & 2 \\ 0 & 0 & 0 & 0 & 0 & 3 \end{pmatrix}, QS_2Q^{-1} = \begin{pmatrix} 1 & 0 & 0 & 1 & 0 & 0 \\ 3 & 3 & 0 & 3 & 0 & 0 \\ 2 & 0 & 3 & 0 & 2 & 0 \\ 3 & 2 & 1 & 3 & 1 & 0 \\ 2 & 0 & 0 & 3 & 2 & 0 \\ 3 & 0 & 2 & 0 & 1 & 3 \end{pmatrix},$$

$$QS_3Q^{-1} = \begin{pmatrix} 0 & 2 & 0 & 2 & 1 & 0 \\ 1 & 3 & 0 & 3 & 1 & 0 \\ 1 & 2 & 1 & 3 & 0 & 0 \\ 3 & 0 & 0 & 3 & 1 & 3 \\ 1 & 2 & 0 & 2 & 1 & 0 \\ 3 & 0 & 0 & 3 & 0 & 0 \end{pmatrix}, QS_4Q^{-1} = \begin{pmatrix} 1 & 2 & 0 & 3 & 2 & 3 \\ 0 & 3 & 0 & 0 & 1 & 1 \\ 2 & 3 & 1 & 2 & 1 & 0 \\ 0 & 2 & 0 & 1 & 1 & 1 \\ 0 & 0 & 2 & 2 & 2 & 1 \\ 0 & 2 & 0 & 2 & 1 & 0 \end{pmatrix}.$$

For $k = 1$ or 4, the matrix QS_kQ^{-1} satisfies $\deg_p((QS_kQ^{-1})_{i,j}) \geqslant s_i - s_j, \forall i > j$. But for $k = 2$ or 3, it does not; for instance, $(QS_2Q^{-1})_{2,1} = 3, (QS_3Q^{-1})_{2,1} = 1$.

We conclude that α_1, α_4 can be lifted, while α_2, α_3 cannot!

References

[1] M. Artin, Algebra, China Mathine Press, Beijing, 2004.

[2] S. F. Du, J. H. Kwak, M. Y. Xu, Linear criteria for lifting automorphisms of elementary abelian regular coverings, Linear Algebra and Its Applica-

tions 373 (2003) 101-119.

[3] J. L. Gross, T. W. Tucker, Generating all graph coverings by permutation voltage assignments, Discrete Mathematics 18 (1977) 273-283.

[4] J. L. Gross, T. W. Tucker, Topological Graph Theory, Wiley-Interscience, New York, 1987.

[5] A. Hatcher, Algebraic Topology, Cambridge University Press, Cambridge, 2002.

[6] B. Kuzman, Arc-transitive elementary abelian covers of the complete graph K_5, Linear Algebra and Its Applications 433 (2010) 1909-1921.

[7] A. Malnič, Group actions, coverings and lifts of automorphisms, Discrete Mathematics 182 (1998) 203-218.

[8] A. Malnič, D. Marušič, P. Potočnik, Elementary abelian covers of graphs, Journal of Algebraic Combinatorics 20 (2004) 71-97.

[9] A. Malnič, P. Potočnik, Invariants subspaces, duality, and covers of the Petersen graph, European Journal of Combinatorics 27 (2006) 971-989.

[10] J-M. Oh, Arc-transitive elementary abelian covers of the Pappus graph, Discrete Mathematics 309 (2009) 6590-6611.